Astronomy & Astrophysics manuscript no. 16471 © ESO 2011
March 1, 2011

arXiv:1101.2044v2 [astro-ph.CO] 28 Feb 2011

Planck Early Results: Statistical properties of extragalactic radio sources in the *Planck* Early Release Compact Source Catalogue

Planck Collaboration: P. A. R. Ade[75], N. Aghanim[51], F. Argüeso[15], M. Arnaud[63], M. Ashdown[61,4], J. Aumont[51], C. Baccigalupi[73], A. Balbi[31], A. J. Banday[79,8,68], R. B. Barreiro[57], J. G. Bartlett[3,59], E. Battaner[80], K. Benabed[52], J.-P. Bernard[79,8], M. Bersanelli[28,45], R. Bhatia[5], A. Bonaldi[41], L. Bonavera[73,6], J. R. Bond[7], J. Borrill[67,76], F. R. Bouchet[52], M. Bucher[3], C. Burigana[44], P. Cabella[31], B. Cappellini[45], J.-F. Cardoso[64,3,52], A. Catalano[3,62], L. Cayón[21], A. Challinor[54,61,11], A. Chamballu[49], R.-R. Chary[50], X. Chen[50], L.-Y Chiang[53], P. R. Christensen[71,32], D. L. Clements[49], S. Colafrancesco[42], S. Colombi[52], F. Couchot[66], B. P. Crill[59,72], F. Cuttaia[44], L. Danese[73], R. D. Davies[60], R. J. Davis[60], P. de Bernardis[27], G. de Gasperis[31], A. de Rosa[44], G. de Zotti[41,73], J. Delabrouille[3], J.-M. Delouis[52], F.-X. Désert[47], C. Dickinson[60], H. Dole[51], S. Donzelli[45,55], O. Doré[59,9], U. Dörl[68], M. Douspis[51], X. Dupac[35], G. Efstathiou[54], T. A. Enßlin[68], H. K. Eriksen[55], F. Finelli[44], O. Forni[79,8], M. Frailis[43], E. Franceschi[44], S. Galeotta[43], K. Ganga[3,50], M. Giard[79,8], G. Giardino[36], Y. Giraud-Héraud[3], J. González-Nuevo[73], K. M. Górski[59,82], S. Gratton[61,54], A. Gregorio[29], A. Gruppuso[44], F. K. Hansen[55], D. Harrison[54,61], S. Henrot-Versillé[66], D. Herranz[57], S. R. Hildebrandt[9,65,56], E. Hivon[52], M. Hobson[4], W. A. Holmes[59], W. Hovest[68], R. J. Hoyland[56], K. M. Huffenberger[81], A. H. Jaffe[49], M. Juvela[20], E. Keihänen[20], R. Keskitalo[59,20], T. S. Kisner[67], R. Kneissl[34,5], L. Knox[23], H. Kurki-Suonio[20,38], G. Lagache[51], A. Lähteenmäki[1,38], A. Lasenby[4,61], R. J. Laureijs[36], C. R. Lawrence[59], S. Leach[73], J. P. Leahy[60], R. Leonardi[35,36,24], P. B. Lilje[55,10], M. Linden-Vørnle[13], M. López-Caniego[57], P. M. Lubin[24], J. F. Macías-Pérez[65], B. Maffei[60], M. Magliocchetti[39], D. Maino[28,45], N. Mandolesi[44], R. Mann[74], M. Maris[43], E. Martínez-González[57], S. Masi[27], M. Massardi[41], S. Matarrese[26], F. Matthai[68], P. Mazzotta[31], P. R. Meinhold[24], A. Melchiorri[27], L. Mendes[35], A. Mennella[28,43], M.-A. Miville-Deschênes[51,7], A. Moneti[52], L. Montier[79,8], G. Morgante[44], D. Mortlock[49], D. Munshi[75,54], A. Murphy[70], P. Naselsky[71,32], P. Natoli[30,2,44], C. B. Netterfield[16], H. U. Nørgaard-Nielsen[13], F. Noviello[51], D. Novikov[49], I. Novikov[71], I. J. O'Dwyer[59], S. Osborne[78], F. Pajot[51], R. Paladini[77,9], B. Partridge[37], F. Pasian[43], G. Patanchon[3], T. J. Pearson[9,50], O. Perdereau[66], L. Perotto[65], F. Perrotta[73], F. Piacentini[27], M. Piat[3], E. Pierpaoli[19], S. Plaszczynski[66], P. Platania[58], E. Pointecouteau[79,8], G. Polenta[2,42], N. Ponthieu[51], T. Poutanen[38,20,1], G. Prézeau[9,59], S. Prunet[52], J.-L. Puget[51], J. P. Rachen[68], R. Rebolo[56,33], M. Reinecke[68], S. Ricciardi[44], T. Riller[68], I. Ristorcelli[79,8], G. Rocha[59,9], C. Rosset[3], M. Rowan-Robinson[49], J. A. Rubiño-Martín[56,33], B. Rusholme[50], A. Sajina[37], M. Sandri[44], D. Scott[18], M. D. Seiffert[59,9], S. Serjeant[17], P. Shellard[11], G. F. Smoot[22,67,3], J.-L. Starck[63,12], F. Stivoli[46], V. Stolyarov[4], R. Stompor[3], R. Sudiwala[75], J.-F. Sygnet[52], J. A. Tauber[36], L. Terenzi[44], L. Toffolatti[14], M. Tomasi[28,45], J.-P. Torre[51], M. Tristram[66], J. Tuovinen[69], M. Türler[48], G. Umana[40], L. Valenziano[44], J. Varis[69], P. Vielva[57], F. Villa[44], N. Vittorio[31], L. A. Wade[59], B. D. Wandelt[52,25], A. Wilkinson[60], D. Yvon[12], A. Zacchei[43], and A. Zonca[24]

(Affiliations can be found after the references)

....

ABSTRACT

The data reported in *Planck*'s Early Release Compact Source Catalogue (ERCSC) are exploited to measure the number counts (dN/dS) of extragalactic radio sources at 30, 44, 70, 100, 143 and 217 GHz. Due to the full-sky nature of the catalogue, this measurement extends to the rarest and brightest sources in the sky. At lower frequencies (30, 44, and 70 GHz) our counts are in very good agreement with estimates based on *WMAP* data, being somewhat deeper at 30 and 70 GHz, and somewhat shallower at 44 GHz. *Planck*'s source counts at 143 and 217 GHz join smoothly with the fainter ones provided by the SPT and ACT surveys over small fractions of the sky. An analysis of source spectra, exploiting *Planck*'s uniquely broad spectral coverage, finds clear evidence of a steepening of the mean spectral index above about 70 GHz. This implies that, at these frequencies, the contamination of the CMB power spectrum by radio sources below the detection limit is significantly lower than previously estimated.

Key words. Surveys - - Radio continuum: general - - Galaxies: active

1. Introduction

Planck[1] (Tauber et al. 2010; Planck Collaboration 2011a) is the third-generation space mission to measure the anisotropy of the cosmic microwave background (CMB). It observes the sky in nine frequency bands covering 30–857 GHz with high sensitivity and angular resolution from $31'$ to $5'$. The Low Frequency Instrument (LFI; Mandolesi et al. 2010; Bersanelli et al. 2010; Mennella et al. 2011) covers the 30, 44, and 70 GHz bands with amplifiers cooled to 20 K. The High Frequency Instrument (HFI; Lamarre et al. 2010; Planck HFI Core Team 2011a) covers the 100, 143, 217, 353, 545, and 857 GHz bands with bolometers cooled to 0.1 K. Polarization is measured in all but the highest two bands (Leahy et al. 2010; Rosset et al. 2010). A combination of radiative cooling and three mechanical coolers produces the temperatures needed for the detectors and optics (Planck Collaboration 2011b). Two data processing centres (DPCs) check and calibrate the data and make maps of the sky (Planck HFI Core Team 2011b; Zacchei et al. 2011). *Planck*'s sensitivity, angular resolution, and frequency coverage make it a powerful instrument for galactic and extragalactic astrophysics as well as cosmology. Early astrophysics results are given in Planck Collaboration, 2011h–z.

---

[1] *Planck* (http://www.esa.int/*Planck*) is a project of the European Space Agency (ESA) with instruments provided by two scientific consortia funded by ESA member states (in particular the lead countries: France and Italy) with contributions from NASA (USA), and telescope reflectors provided in a collaboration between ESA and a scientific consortium led and funded by Denmark.

The *Planck* Early Release Compact Source Catalogue (ERCSC, Planck Collaboration (2011c)) reports data on sources detected during the first 1.6 full-sky surveys, and thus offers, among other things, the opportunity of studying the statistical properties of extragalactic sources over a broad frequency range never fully explored by blind surveys. We will focus here on counts of extragalactic radio sources and on their spectral properties in the 30–217 GHz range.[2]

Although knowledge of the statistical properties at high radio frequency for this population of extragalactic sources has greatly improved in the recent past – thanks to many ground-based observational campaigns and to the Wilkinson Microwave Anisotropy Probe (*WMAP*) surveys from space – above about 70 GHz these properties are still largely unknown or very uncertain. This is essentially due to the fact that very large area surveys at mm wavelengths are made difficult by the small fields of view of ground-based radio telescopes and by the long integration times required.

The most recent estimates on source number counts up to $\sim 50-70$ GHz, and the optical identifications of the corresponding bright point sources (see, e.g., Massardi et al. (2008, 2010)), show that these counts are dominated by radio sources whose average spectral index is "flat", i.e., $\alpha \simeq 0.0$ (with the usual convention $S_\nu \propto \nu^\alpha$). This result confirms that the underlying source population is essentially made of Flat Spectrum Radio Quasars (FSRQ) and BL Lac objects, collectively called blazars,[3] with minor contributions coming from other source populations (Toffolatti et al. 1998; de Zotti et al. 2005). At frequencies > 100 GHz, however, there is now new information for sources with flux densities below about 1 Jy coming from the South Pole Telescope (SPT) collaboration (Vieira et al. 2010), with surveys over 87 deg^2 at 150 and 220 GHz, and from the Atacama Cosmology Telescope (ACT) survey over 455 deg^2 at 148 GHz (Marriage et al. 2010).

The "flat" spectra of blazars are generally believed to result from the superposition of different components in the inner part of AGN relativistic jets, each with a different synchrotron self-absorption frequency (Kellermann & Pauliny-Toth 1969). At a given frequency, the observed flux density is thus dominated by the synchrotron-emitting component which becomes self-absorbed and, in the equipartition regime, the resulting spectrum is approximately flat. However, this "flat" spectrum cannot be maintained up to very high frequencies, because of electron energy losses in the dominant jet-emission component (i.e., electron ageing), or the transition to the optically-thin regime, with the onset of a "steep" spectrum with a standard spectral index $\alpha = -0.7$ to -0.8. A slightly steepened spectrum may also be caused by the superposition of many jet components. The redshift moves the observed steepening to lower frequencies and, thus, a greater fraction of blazar sources are observed with a steep spectrum at sub-mm wavelengths. With current data it is not yet possible to decide among the different scenarios. However, given their sensitivity and full sky coverage, *Planck* surveys are uniquely able to shed light on this transition from an almost "flat" to a "steep" regime in the spectra of blazar sources.

The outline of this paper is as follows. In § 2.1 we briefly sketch the main properties of the ERCSC. In § 2.2 we summarize the source validation. In § 2.3 we describe the complete sample selected at 30 GHz, used for the analysis of spectral properties. In § 3 we present the source counts over the frequency range 30–217 GHz. In § 4 we investigate the spectral index distributions in different frequency intervals. Finally, in § 5 we summarize our main conclusions.

2. The *Planck* ERCSC

2.1. Overview

The *Planck* ERCSC (Planck Collaboration 2011c) lists positions and flux densities for the compact sources recovered from the *Planck* first 1.6 full sky survey maps in nine frequency channels between 30 and 857 GHz. Thus about 60% of the sources have been covered twice, with a time separation of about 6 months. Sources near the ecliptic poles, where the scan circles intersect, are often covered multiple times. ERCSC flux densities are therefore averages over different observing time periods. They have been calculated by aperture photometry, using the most recent definition of the beam shapes and sizes (Planck HFI Core Team 2011b; Zacchei et al. 2011). In the frequency range considered in this paper, the *Planck* photometric calibration is based on the CMB dipole and on the modulation induced on it by the spacecraft orbital motion. According to Zacchei et al. (2011) the absolute photometric calibration at LFI frequencies (30, 44 and 70 GHz) is at the $\sim 1\%$ level, while Planck HFI Core Team (2011b) reports on a relative photometric accuracy, between the frequency channels from 100 to 353 GHz, better than 2% and, more likely, at the $\sim 1\%$ level.

---

[2] In all our calculations we have used the effective central frequencies for the *Planck* channels (Mennella et al. 2011; Planck HFI Core Team 2011a), although we indicate their nominal values. The most relevant difference is at 30 GHz, with a central frequency of 28.5.

[3] Blazars are jet-dominated extragalactic objects characterized by a strongly variable and polarized emission of the nonthermal radiation, from low radio energies up to the high energy gamma rays (Urry & Padovani 1995). Detailed analyses of Spectral Energy Distributions (SEDs) of complete blazar samples built by using simultaneous *Planck* , Swift and Fermi data are given in (Planck Collaboration 2011k).



Fig. 1. Comparison between the ERCSC flux densities at 30 GHz (left panel) and at 44 GHz (right panel) with the almost simultaneous ATCA measurements (PACO project) at 32.2 and 39.7 GHz, respectively. No correction for the slightly different frequencies has been applied.

The final version of the ERCSC (see also Planck Collaboration 2011c for more details) has been created by specifically applying the so-called "PowellSnakes" detection method (Carvalho et al. 2009) to the *Planck* full sky (dipole subtracted) anisotropy maps in the frequency channels from 30 to 143 GHz and the SExtractor package (Bertin & Arnouts 1996) in the four channels at higher frequencies (from 217 to 857 GHz). A "compact" source is accepted in the ERCSC if it survived a set of primary and secondary selection criteria. The primary criterion utilized the feedback from the Monte Carlo Quality Assessment system and introduced a signal-to-noise ratio cut to ensure that $> 90\%$ of the sources in the catalogue are reliable and have a flux density accuracy better than 30%. The secondary criterion comprised a set of cuts that removed the extended sources with an elongation > 3.0 (the ratio between the major and minor axis, in pixels, of the detected source) and sources that could be potentially spurious. For instance a source is dropped whenever $> 5\%$ of the pixels within $2\times$FWHM of its position had invalid values (see the ERCSC Explanatory Supplement for more details). The overriding requirement in constructing the ERCSC is source reliability and not completeness.

2.2. Validation and photometry check of ERCSC sources

The validation process for the compact sources included in the *Planck* ERCSC was performed by two different teams, selected among members of the *Planck* Consortia: a Validation Team on radio sources (VTRS) and a Validation Team on far-IR sources. The two teams worked separately at first, but cross-checked their results in the second phase of the process. The processing steps and main outcomes are summarized in the Explanatory Supplement released with the ERCSC. For compact radio (i.e., synchrotron dominated) sources, the VTRS (see Planck Collaboration (2011j) for a more detailed discussion) has found that $> 97\%$ of the ERCSC sources at 30 GHz have reliable counterparts in published catalogues at GHz frequencies (PMN: Wright et al. 1996; GB6: Gregory et al. 1996; NVSS: Condon et al. 1998; SUMSS: Mauch et al. 2003; AT20G: Massardi et al. 2008; Murphy et al. 2010). Similar (although slightly lower) percentages were found for ERCSC sources detected at 44 and 70 GHz. At higher frequencies (≥ 100 GHz) *Planck* detects an increasing fraction of dusty galaxies, undetected by low-frequency surveys. Therefore, the source reliability was confirmed by internal matches of sources detected in two neighboring *Planck* frequency channels: i.e., 143 and 217 GHz, or 217 and 353 GHz, etc.. However, the validation of synchrotron-dominated sources is still relatively easy to perform, since all of them must be present in low-frequency catalogues.

The *WMAP* 7-year catalogue (Gold et al. 2010) contains a total of 471 sources detected in at least one frequency channel. Of these, 289, 281, 166 and 59 sources are detected as $\geq 5\sigma$ peaks in the 33, 41, 61, and 94 GHz maps, respectively. The ERCSC catalogue includes 88%, 63%, 81%, and 95% of the 5σ *WMAP* sources at 30, 44, 70, and 100 GHz,respectively. The median of the distribution of offsets between *WMAP* and *Planck* positions at each frequency are $2.5'$, $2.1'$, $1.7'$, and $1.0'$ at each frequency (see also Planck Collaboration 2011j for a more detailed discussion on this subject). Except for the 44 GHz channel, where *Planck* is known to be less sensitive, most *WMAP* sources that failed to be included in the *Planck* ERCSC (31 sources at 30 GHz) are generally at the faint end of the flux density distribution (i.e., near the detection threshold) and may have flux densities boosted by the Eddington bias or the effects of confusion, or may be spurious. The absence from the ERCSC of a few brighter *WMAP* sources (5 sources at 30 GHz) is probably caused by their variability.

The *Planck*-ATCA Co-eval Observations (PACO) project (Massardi et al. 2011) has provided measurements with the Australia Telescope Compact Array (ATCA) of sources potentially detectable by *Planck* almost simultaneously with *Planck* observations. 147 ERCSC sources have PACO observations at 32.2 and 39.7 GHz within 10 days of *Planck* observations. All these sources are unresolved also by ATCA. As illustrated in Fig. 1, the comparison between ATCA and ERCSC flux densities at the nearest frequen-



Fig. 2. Euclidean normalized differential number counts at the LFI frequencies. The red circles with Poisson error bars show the counts of sources with counterparts in our reference 30 GHz sample. In each panel, the solid curves show the total number counts of extragalactic radio sources predicted by the de Zotti et al. (2005) model. Also shown are: the counts estimated at 31 GHz from DASI (grey dashed box; Kovac et al. (2002)) and at 33 GHz from the VSA data (grey box; Cleary et al. (2005)); the counts from the PACO (grey diamonds; Bonavera et al., in preparation) and the *WMAP* 5-yr surveys (grey squares; Massardi et al. (2009)), at the closest frequencies, and the counts estimated by Waldram et al. (2007) (grey dashed line), exploiting multi-frequency follow-up observations of the 15 GHz 9C sources. The vertical dashed magenta line in the upper panel indicates the flux density completeness limit, 1.0 Jy, estimated for our primary sample § 2.3.

cies (30 and 44 GHz, respectively), shows a reassuringly close agreement. The faintest ERCSC flux densities are obviously enhanced by the effect of the Eddington bias,the noise-increased number count of point sources to a given detection threshold, which is besides enhanced in the 44 GHz *Planck* channel, where the noise level is higher.

2.3. The 30 GHz extragalactic radio source sample

2.3.1. Identification of compact Galactic sources

To minimize the contamination of the sample by Galactic sources we have restricted ourselves to $|b| > 5^\circ$ and we have also excluded sources within 5° and 2.5°, respectively, of the nominal centres of the Large and Small Magellanic Clouds. Outside these regions, a search in the SIMBAD database, with a search radius of $16'$, corresponding to about half the FWHM at 30 GHz, has yielded 18 associations of ERCSC sources with known Galactic objects (5 PNe, 10 HII regions, and 3 SNRs), all within $5'$ of the ERCSC position. After having removed these sources we are left with 533 compact extragalactic radio sources detected at 30 GHz, with 97% or more of them identified in external catalogues at GHz frequencies. This constitutes our primary sample.

2.3.2. Completeness and uniformity tests

An indication of the completeness limit of our sample is obtained by looking at the differential counts (see top panel in Fig. 2) : a sharp decrease of the slope at faint flux densities ($S_\nu \lesssim 1$ Jy at 30 GHz) signals the onset of incompleteness. Based on the ATCA 20 GHz counts (Massardi et al. 2008; Murphy et al. 2010), we expect that the slope of the counts remains approximately constant over the limited flux density range covered by the ERCSC. Therefore, the flux density interval (or bin size) containing a fixed number of sources must decrease as a power of the central flux density. As shown in the top panel of Fig. 3, at 30 GHz this happens down to a flux density of about 0.9 Jy, where the curve abruptly flattens.

We need also to test whether the spatial distribution of sources in our sample is consistent with being statistically uniform, as it must be in the case of extragalactic sources. Deviations from uniformity may be expected at lower Galactic latitudes, both because of residual contamination by unrecognized Galactic sources and through the effect of a stronger Eddington bias due to fluctuations of diffuse Galactic emission. The bottom panel of Fig. 3 shows no significant deviations from a uniform distribution on the sky for $|b| > 5°$ if we adopt a completeness limit of $\simeq 1.0$ Jy at 30 GHz. Remarkably, the average source density at $|b| > 5°$, $D = 24.23$ (in sources per sr), is very similar to the value found at $|b| > 30°$, $D = 23.71$, which guarantees that we are not losing extragalactic sources, in this frequency range, when going down to lower Galactic latitudes, and that the residual contamination due to unrecognized Galactic sources is negligible. Larger deviations from uniformity are found if we adopt fainter completeness limits. Taking into account both results, we therefore adopt S_ν=1.0 Jy as an estimate for the completeness limit at 30 GHz. Our primary sample is made of 290 sources above the adopted $S_\nu = 1.0$ Jy flux density limit. As a comparison, Massardi et al. (2009) detected 281 sources at $|b| > 5^\circ$ and $S_\nu \gtrsim 1$ Jy in their blind survey performed on the 5-year *WMAP* 33 GHz map.

In addition, we used the *Planck* HFI frequency channels at 143 and 217 GHz to select all the extragalactic sources in the ERCSC whose spectra are still dominated by non-thermal synchrotron emission at 217 GHz (409 sources with $\alpha_{143}^{217} < 0.5$; see Fig. 4). We limit the selection of this secondary sample to the above α_{143}^{217} value with the purpose of excluding all possible sample contamination coming from a second population of sources dominated by thermal dust emission ($\alpha_{143}^{217} > 1.0$). The choice of $\alpha_{143}^{217} < 1.0$, corresponding to a minimum in the distribution, would not change our results since very few sources (< 10) have spectral indices in the interval $0.5 < \alpha_{143}^{217} < 1.0$. This selection (at above 100 GHz, where there is not yet a complete match of ERCSC sources with external catalogs) is useful for comparing the outcomes from this (secondary) sample with our predictions on the statistical properties of extragalactic sources in our primary sample, selected at 30 GHz.

3. Number counts

Figures 2 and 5 show the number counts of extragalactic radio sources at the six *Planck* frequencies from 30 to 217 GHz (see also Table 1). The sharp breaks in the number counts at approximately 1.0 Jy (30 GHz), 1.5 Jy (44 GHz), 1.1 Jy (70 GHz), 0.9 Jy (100 GHz), 0.5 Jy (143 GHz) and 0.4 Jy (217 GHz) signal the onset of incompleteness.

The results of deeper surveys on small fractions of the sky and the *WMAP* differential number counts are also shown, for comparison. The agreement with *WMAP* data is very good. Our differential counts at 30 and 70 GHz are



Fig. 3. Completeness and uniformity tests at 30 GHz. Top panel: flux density interval, or bin size, including a fixed number of sources as a function of flux density. The change of slope at about 0.9 Jy signals the onset of incompleteness. Bottom panel: uniformity test for sources with flux densities $S_\nu \geq 1.0$ Jy. The source density, D, in sources per sr, within regions at different Galactic latitudes, shows an acceptable uniformity. The horizontal grey (solid) and blue (dashed) lines show the average source density at Galactic latitudes above $|b| = 5$ and $|b| = 30$ deg, respectively. As it is apparent, the two values are very close to each other, within the 1σ level calculated for the overall population in the two sky areas considered here, and well inside the 1σ normalized Poisson error bars. We also checked that at flux densities below about 0.9–1.0 Jy statistically more relevant deviations from a uniform distribution begin to appear.



Fig. 4. Spectral index distribution of ERCSC sources between 143 and 217GHz. Only sources detected at 143 GHz and at 217 GHz have been considered: no upper limits on flux densities have been used in this calculation.

Table 1. Euclidean-normalized differential number counts per steradian estimated from the *Planck* ERCSC at 30–217 GHz. The 143 and 217 GHz number counts are those represented by blue diamonds in Fig. 5. The bins are centred in the $\log_{10}(S_\nu)$ values and are symmetric in logarithmic scale.

	$S^{5/2}dN/dS$ [Jy$^{3/2}$sr^{-1}]					
$\log_{10}(S_\nu)$ [Jy]	30 [GHz]	44 [GHz]	70 [GHz]	100 [GHz]	143 [GHz]	217 [GHz]
−0.5	...	...	...	...	0.9 ± 0.2	4.6 ± 0.6
−0.4	...	...	...	1.1 ± 0.4	6.3 ± 0.8	8.2 ± 0.9
−0.3	2.6 ± 0.6	...	...	5.8 ± 0.9	11.6 ± 1.2	8.9 ± 1.1
−0.2	9.5 ± 1.3	...	1.3 ± 0.6	13.3 ± 1.6	11.6 ± 1.5	8.5 ± 1.3
−0.1	29 ± 3	2.1 ± 0.7	7.0 ± 1.4	20 ± 2	12.6 ± 1.8	10.5 ± 1.7
0.0	48 ± 4	10 ± 2	23 ± 3	25 ± 3	16 ± 2	11 ± 2
0.1	42 ± 5	25 ± 4	29 ± 4	22 ± 3	12 ± 2	14 ± 3
0.2	25 ± 4	36 ± 5	16 ± 3	18 ± 4	15 ± 3	7 ± 2
0.3	38 ± 6	35 ± 6	24 ± 5	18 ± 4	20 ± 5	12 ± 4
0.4	41 ± 8	50 ± 9	23 ± 6	21 ± 6	11 ± 4	6 ± 3
0.5	34 ± 8	17 ± 6	28 ± 8	15 ± 6	11 ± 5	13 ± 5
0.6	30 ± 9	27 ± 9	33 ± 10	21 ± 9	12 ± 6	9 ± 5
0.7	47 ± 14	38 ± 12	9 ± 6	4 ± 4	13 ± 7	13 ± 7
0.8	18 ± 10	6 ± 5	12 ± 8	18 ± 10	12 ± 8	12 ± 8
0.9	34 ± 16	34 ± 16	8 ± 7	17 ± 11	8 ± 7	8 ± 7
1.0	24 ± 15	...	36 ± 20	36 ± 20	24 ± 15	...

somewhat deeper than the *WMAP* ones at 33 and 61 GHz, while at 44 GHz they are somewhat shallower than the ones calculated for the 41 GHz *WMAP* channel. Also our counts above the completeness limits appear to join smoothly with those from deeper surveys.

At frequencies of up to 100 GHz, the predictions of the de Zotti et al. (2005) cosmological evolution model – relying on extrapolations from lower frequency data and capable of providing a very good fit to almost all data on number counts as well as on other statistics of radio sources at frequencies above 5 GHz – are in generally good agreement with our current findings. This result implies that no new radio source population shows up at bright flux densities. Very few "extreme" or "inverted–spectrum" compact radio sources are found in the *Planck* ERCSC. The emission and spectral properties of these sources, which are interesting in their own right, are discussed in a companion paper (Planck Collaboration 2011j).

At higher frequencies (i.e., at 143 and 217 GHz) we also plotted the number counts obtained by using the sample of radio sources selected at 143 and 217 GHz (blue diamonds; see § 2.3 and Fig. 4 for more details). These number counts (calculated from our secondary sample, 284 sources at above 0.5 Jy) turn out to be in almost perfect agreement with the ones obtained in the previous Section from our primary sample selected at 30 GHz (290 sources at $S_\nu \geq 1.0$), thus confirming that no bright extragalactic radio sources are missed by our selection criteria and that the underlying parent population turns out to be (statistically) the same.

Figure 5 shows that the de Zotti et al. (2005) model over-predicts the bright counts by a factor of about 2 at 143 GHz and about 2.6 at 217 GHz, while it is consistent with the fainter SPT (Vieira et al. 2010) and ACT (Marriage et al. 2010) counts. As discussed in the next Section, the discrepancy between the model and our current data is due to a steepening of the spectra of ERCSC sources above about 70 GHz, not predicted by the model but, at least partially, already suggested by other data sets (González-Nuevo et al. 2008; Sadler et al. 2008).

An implication of this result is that the contamination of the CMB angular power spectrum by extragalactic radio sources below the detection limit at 143 and 217 GHz is lower than predicted by the de Zotti et al. (2005) model. Assuming a Poisson distribution (clustering effects are reduced to negligible values by the very broad luminosity function of radio sources, e.g., Toffolatti et al. (2005); de Zotti et al. (2010)) and simply scaling down the model counts by the factors mentioned above, the amplitude of the angular power spectrum of unresolved sources goes down by roughly the same factor [4]. This is, however, an upper limit to the correction factor, especially at 217 GHz, because if we apply the factor calculated above to all flux densities we would end up with a clear underestimate of the faint counts measured by Vieira et al. (2010). Therefore, if we consider much fainter source detection limits, as foreseen for future experiments in the sub-mm, the amplitude of the angular power spectrum due to unresolved sources stays essentially at the same level as predicted by the De Zotti et al. model.

4. Spectral index distributions

To study the spectral properties of the extragalactic radio sources in the *Planck* ERCSC we used our reference 30 GHz sample above the estimated completeness limit (1.0 Jy; § 2.3). Not all of these sources were detected at the $\geq 5\sigma$ level in each of the *Planck* frequency channels considered. Whenever a source was not detected in a given channel we replaced its (unknown) flux density by a 5σ upper limit, where for σ we used the average r.m.s. error estimated at each *Planck* frequency. The upper limits have been redistributed among the flux density bins by using a Survival

---

[4] As an example, the amplitude of the power spectrum of radio sources, in terms of C_ℓ values, below, e.g., $S_\nu \leq 1$ Jy at 217 GHz, corresponds to $\sim 4.2 \times 10^{-5} \mu\mathrm{K}^2$, if calculated from the *Planck* ERCSC differential number counts of Table 1. For comparison, undetected radio sources sum up $\sim 1.1 \times 10^{-4} \mu\mathrm{K}^2$, if we integrate the differential number counts of the de Zotti et al. (2005) cosmological evolution model up to $S_\nu = 1$ Jy.



Fig. 5. Euclidean normalized differential number counts at the HFI frequencies (100, 143, and 217 GHz). The red circles with Poisson error bars show the counts of sources with counterparts in our reference 30 GHz sample. At 143 and 217 GHz the blue diamonds (shifted to the left by half of the bin size, for clarity) show the counts obtained after removing sources with 143–217 GHz spectral index indicative of dust emission (see Sect. 2.3.2). Again, in each panel, the solid curves show the total number counts of extragalactic radio sources predicted by the de Zotti et al. (2005) evolution model. Also shown are the SPT (grey squares; Vieira et al. 2010) and ACT (grey stars; Marriage et al. 2010 counts of radio sources. At 100 GHz we also show the estimated counts by Sadler et al. (2008) from follow-up observations of a sample of sources selected from the 20 GHz ATCA survey (grey dashed line).

Table 2. Median and standard deviations of the spectral index distributions between 30 GHz and the selected frequency. We adopt the convention $S_\nu \propto \nu^\alpha$.

ν[GHz]	44	70	100	143	217
median	-0.06	-0.18	-0.28	-0.39	-0.37
error	0.01	0.01	0.01	0.01	0.01
σ	0.30	0.18	0.17	0.16	0.15

Analysis technique and, more specifically, by adopting the Kaplan-Meyer estimator (as implemented in the ASURV code, (Lavalley et al. 1992)). Since the fraction of upper limits is always small (it reaches approximately 30% only in our less sensitive channel at 44GHz), the spectral index distributions are reliably reconstructed at each frequency.

Table 2 gives the median spectral indices between 30 GHz and the other frequencies considered here and the dispersions of the spectral index distributions. A moderate steepening of spectral indices at higher frequencies is clearly apparent. Hints in this direction were previously found by González-Nuevo et al. (2008) from their analysis of the NEWPS sample (López-Caniego et al. 2007) and also by Sadler et al. (2008). Additional evidence of spectral steepening is presented in Planck Collaboration (2011k).

Our 30–100 GHz spectral index is close to the $\alpha \simeq -0.39$ found by Sadler et al. (2008) between 20 and 95 GHz, for a sample with 20 GHz flux density $S > 150$ mJy. Moreover, our 30–143 GHz median spectral index is in very good agreement with the one found by Marriage et al. (2010) for their bright ($S_\nu > 50$ mJy) 148 GHz-selected sample with



Fig. 6. Spectral index distributions for different frequency intervals calculated by taking into account all sources selected at 30 GHz with $S_\nu > 1$ Jy. There is clear evidence for a steepening above 70 GHz.



Fig. 7. Contour levels of the distribution of α_{70}^{143} vs. α_{30}^{70} obtained by Survival Analysis, i.e., taking into account the upper limits to flux densities at each frequency. The colour scale can be interpreted as the probability of having any particular pair of values of the two spectral indices. The maximum probability corresponds to $\alpha_{30}^{70} \simeq -0.18$ and $\alpha_{70}^{143} \simeq -0.50$.

complete cross-identifications from the Australia Telescope 20 GHz survey, i.e $\alpha_{20}^{148} = -0.39 \pm 0.04$. On the other hand, (Vieira et al. 2010) find that their much fainter synchrotron emitting radio sources selected at 150 GHz are consistent with a flatter spectral behaviour (mean $\alpha_5^{150} \simeq -0.1$) between 5 GHz and 150 GHz. Massardi et al. (2010) find mean spectral indices $\alpha_5^{150} \simeq -0.17$ and $\alpha_{20}^{150} \simeq -0.30$ for AT20G sources with 150 GHz flux density $S > 50$ mJy. A flattening of the mean/median high-frequency spectral indices at flux densities fainter than the ones probed by the *Planck* ERCSC may help to account for the unusually "flat" normalized counts at 143 and 217 GHz.

In Fig. 6 we compare the distributions of spectral indices over different frequency intervals. There is a clear shift toward steeper values above 70 GHz: the median values vary from $\alpha_{30}^{70} = -0.18 \pm 0.01$ ($\sigma = 0.18$) to $\alpha_{70}^{143} = -0.52 \pm 0.01$ ($\sigma = 0.22$). On the other hand, the distribution of spectral indices between 143 and 217 GHz is close to the one found for α_{70}^{143} ($\alpha_{143}^{217} = -0.46 \pm 0.01$; $\sigma = 0.23$). This latter result is again very similar to the corresponding value calculated for all the sources detected at 143 and 217 GHz with $\alpha_{143}^{217} < 0.5$, i.e., $\alpha_{143}^{217} = -0.51 \pm 0.01$, as shown in Fig. 4. Moreover, in the paper Planck Collaboration (2011k) an average value of $\alpha = -0.56$ ($\sigma = 0.29$) at HFI frequencies is found for their sample of 84 bright blazars selected at 37 GHz, with flux densities measured in at least 3 HFI channels, in full agreement with our present findings.

Figure 7 presents the contour levels of the distribution of α_{70}^{143} vs. α_{30}^{70} (obtained using Survival Analysis) in the form of a 2D probability field: the colour scale can be interpreted as the probability of a given pair of spectral indices.

As already noted, at high *Planck* frequencies most of the extragalactic radiosources are blazars. From the contour plot of Fig. 7 it is possible to see that the maximum probability corresponds to $\alpha_{30}^{70} \simeq -0.18$ and $\alpha_{70}^{143} \simeq -0.5$. A secondary maximum can also be seen at $\alpha_{70}^{143} \simeq -1.2$. However, a physical interpretation of these features goes beyond the purposes of this work and, moreover, more data at higher frequencies are needed. A detailed discussion on the modelling of the spectra of this source population is presented in a companion paper (Planck Collaboration 2011k).

5. Conclusions

We have exploited the ERCSC to estimate the bright counts of extragalactic radio sources at 6 frequencies (30, 44, 70, 100, 143, and 217 GHz) and to investigate the spectral properties of sources in a complete sample selected at 30 GHz. The counts at 30, 44, and 70 GHz are in good agreement with those derived from *WMAP* data at nearby frequencies. The completeness limit of the ERCSC is somewhat deeper than that of *WMAP* at 30 and 70 GHz and somewhat shallower at 44 GHz. At higher frequencies the ERCSC has allowed us to obtain the first estimate of the differential number counts at bright flux density levels. At 30, 143 and 217 GHz, the present counts join smoothly to those from deeper surveys over small fractions of the sky.

The de Zotti et al. (2005) model is consistent with the present counts at frequencies up to 100 GHz, but overpredicts the counts at higher frequencies by a factor of about 2.0 at 143 GHz and about 2.6 at 217 GHz. This implies that the contamination of the CMB power spectrum by radio sources below the 1 Jy detection limit is lower than previously estimated. No significant changes are found, however, if we consider fainter source detection limits, i.e., 100 mJy, given the convergence between predicted and observed number counts.

The analysis of the spectral index distribution over different frequency intervals, within the uniquely broad range covered by *Planck* in the mm and sub-mm domain, has highlighted an average *steepening* of source spectra above about 70 GHz. The median values of spectral indices vary from $\alpha_{30}^{70} = -0.18 \pm 0.01$ ($\sigma = 0.18$) to $\alpha_{70}^{143} = -0.52 \pm 0.01$ ($\sigma = 0.22$). This steepening accounts for the discrepancy between the de Zotti et al. (2005) model predictions and the observed differential number counts at HFI frequencies. The current outcome is also in agreement with the findings of Planck Collaboration (2011k) on a complete sample of blazars selected at 37 GHz. The change detected in the spectral behaviour of extragalactic radio sources in the *Planck* ERCSC at frequencies above 70-100 GHz can be tentatively explained by electron ageing or by the transition to the optically thin regime, predicted in current models for radio emission in blazar sources. However, with present data it is not yet possible to clarify the situation. In the near future, the data of the *Planck* Legacy Survey will surely prove very useful in settling this open issue.

Acknowledgements. The *Planck* Collaboration thanks the referee, Ronald Ekers, for his insightful comments, that helped improving the paper. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. The Planck Collaboration acknowledges the support of: ESA; CNES and CNRS/INSU-IN2P3-INP (France); ASI, CNR, and INAF (Italy); NASA and DoE (USA); STFC and UKSA (UK); CSIC, MICINN and JA (Spain); Tekes, AoF and CSC (Finland); DLR and MPG (Germany); CSA (Canada); DTU Space (Denmark); SER/SSO (Switzerland); RCN (Norway); SFI (Ireland); FCT/MCTES (Portugal); and DEISA (EU). A description of the Planck Collaboration and a list of its members can be found at `http://www.rssd.esa.int/index.php?project=PLANCK&page=Planck_Collaboration`

References

Bersanelli, M., Mandolesi, N., Butler, R. C., et al. 2010, A&A, 520, A4+
Bertin, E. & Arnouts, S. 1996, A&AS, 117, 393
Carvalho, P., Rocha, G., & Hobson, M. P. 2009, MNRAS, 393, 681
Cleary, K. A., Taylor, A. C., Waldram, E., et al. 2005, MNRAS, 360, 340
Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, AJ, 115, 1693
de Zotti, G., Massardi, M., Negrello, M., & Wall, J. 2010, A&A Rev., 18, 1
de Zotti, G., Ricci, R., Mesa, D., et al. 2005, A&A, 431, 893
Gold, B., Odegard, N., Weiland, J. L., et al. 2010, ArXiv e-prints
González-Nuevo, J., Massardi, M., Argüeso, F., et al. 2008, MNRAS, 384, 711
Gregory, P. C., Scott, W. K., Douglas, K., & Condon, J. J. 1996, ApJS, 103, 427
Kellermann, K. I. & Pauliny-Toth, I. I. K. 1969, ApJ, 155, L71+
Kovac, J. M., Leitch, E. M., Pryke, C., et al. 2002, Nature, 420, 772
Lamarre, J., Puget, J., Ade, P. A. R., et al. 2010, A&A, 520, A9+
Lavalley, M. P., Isobe, T., & Feigelson, E. D. 1992, in Bulletin of the American Astronomical Society, Vol. 24, Bulletin of the American Astronomical Society, 839–840
Leahy, J. P., Bersanelli, M., D'Arcangelo, O., et al. 2010, A&A, 520, A8+
López-Caniego, M., González-Nuevo, J., Herranz, D., et al. 2007, ApJS, 170, 108
Mandolesi, N., Bersanelli, M., Butler, R. C., et al. 2010, A&A, 520, A3+
Marriage, T. A., Juin, J. B., Lin, Y., et al. 2010, ArXiv e-prints
Massardi, M., Bonaldi, A., Bonavera, L., et al. 2011, ArXiv e-prints
Massardi, M., Bonaldi, A., Negrello, M., et al. 2010, MNRAS, 404, 532
Massardi, M., Ekers, R. D., Murphy, T., et al. 2008, MNRAS, 384, 775
Massardi, M., López-Caniego, M., González-Nuevo, J., et al. 2009, MNRAS, 392, 733
Mauch, T., Murphy, T., Buttery, H. J., et al. 2003, MNRAS, 342, 1117
Mennella et al. 2011, Planck early results 03: First assessment of the Low Frequency Instrument in-flight performance (Submitted to A&A)
Murphy, T., Sadler, E. M., Ekers, R. D., et al. 2010, MNRAS, 402, 2403
Planck Collaboration. 2011a, Planck early results 01: The Planck mission (Submitted to A&A)
Planck Collaboration. 2011b, Planck early results 02: The thermal performance of Planck (Submitted to A&A)
Planck Collaboration. 2011c, Planck early results 07: The Early Release Compact Source Catalogue (Submitted to A&A)
Planck Collaboration. 2011d, Planck early results 08: The all-sky early Sunyaev-Zeldovich cluster sample (Submitted to A&A)
Planck Collaboration. 2011e, Planck early results 09: XMM-Newton follow-up for validation of Planck cluster candidates (Submitted to A&A)
Planck Collaboration. 2011f, Planck early results 10: Statistical analysis of Sunyaev-Zeldovich scaling relations for X-ray galaxy clusters (Submitted to A&A)
Planck Collaboration. 2011g, Planck early results 11: Calibration of the local galaxy cluster Sunyaev-Zeldovich scaling relations (Submitted to A&A)
Planck Collaboration. 2011h, Planck early results 12: Cluster Sunyaev-Zeldovich optical Scaling relations (Submitted to A&A)
Planck Collaboration. 2011i, Planck early results 13: Statistical properties of extragalactic radio sources in the Planck Early Release Compact Source Catalogue (Submitted to A&A)
Planck Collaboration. 2011j, Planck early results 14: Early Release Compact Source Catalogue validation and extreme radio sources (Submitted to A&A)
Planck Collaboration. 2011k, Planck early results 15: Spectral energy distributions and radio continuum spectra of northern extragalactic radio sources (Submitted to A&A)
Planck Collaboration. 2011l, Planck early results 16: The Planck view of nearby galaxies (Submitted to A&A)
Planck Collaboration. 2011m, Planck early results 17: Origin of the submillimetre excess dust emission in the Magellanic Clouds (Submitted to A&A)
Planck Collaboration. 2011n, Planck early results 18: The power spectrum of cosmic infrared background anisotropies (Submitted to A&A)
Planck Collaboration. 2011o, Planck early results 19: All-sky temperature and dust optical depth from Planck and IRAS — constraints on the "dark gas" in our Galaxy (Submitted to A&A)
Planck Collaboration. 2011p, Planck early results 20: New light on anomalous microwave emission from spinning dust grains (Submitted to A&A)
Planck Collaboration. 2011q, Planck early results 21: Properties of the interstellar medium in the Galactic plane (Submitted to A&A)
Planck Collaboration. 2011r, Planck early results 22: The submillimetre properties of a sample of Galactic cold clumps (Submitted to A&A)
Planck Collaboration. 2011s, Planck early results 23: The Galactic cold core population revealed by the first all-sky survey (Submitted to A&A)
Planck Collaboration. 2011t, Planck early results 24: Dust in the diffuse interstellar medium and the Galactic halo (Submitted to A&A)
Planck Collaboration. 2011u, Planck early results 25: Thermal dust in nearby molecular clouds (Submitted to A&A)
Planck Collaboration. 2011v, The Explanatory Supplement to the Planck Early Release Compact Source Catalogue (ESA)
Planck HFI Core Team. 2011a, Planck early results 04: First assessment of the High Frequency Instrument in-flight performance (Submitted to A&A)
Planck HFI Core Team. 2011b, Planck early results 06: The High Frequency Instrument data processing (Submitted to A&A)
Rosset, C., Tristram, M., Ponthieu, N., et al. 2010, A&A, 520, A13+
Sadler, E. M., Ricci, R., Ekers, R. D., et al. 2008, MNRAS, 385, 1656
Tauber, J. A., Mandolesi, N., Puget, J., et al. 2010, A&A, 520, A1+
Toffolatti, L., Argueso Gomez, F., de Zotti, G., et al. 1998, MNRAS, 297, 117
Toffolatti, L., Negrello, M., González-Nuevo, J., et al. 2005, A&A, 438, 475
Urry, C. M. & Padovani, P. 1995, PASP, 107, 803
Vieira, J. D., Crawford, T. M., Switzer, E. R., et al. 2010, ApJ, 719, 763
Waldram, E. M., Bolton, R. C., Pooley, G. G., & Riley, J. M. 2007, MNRAS, 379, 1442
Wright, A. E., Griffith, M. R., Hunt, A. J., et al. 1996, ApJS, 103, 145
Zacchei et al. 2011, Planck early results 05: The Low Frequency Instrument data processing (Submitted to A&A)

---

[1] Aalto University Metsähovi Radio Observatory, Metsähovintie 114, FIN-02540 Kylmälä, Finland

[2] Agenzia Spaziale Italiana Science Data Center, c/o ESRIN, via Galileo Galilei, Frascati, Italy

[3] Astroparticule et Cosmologie, CNRS (UMR7164), Université Denis Diderot Paris 7, Bâtiment Condorcet, 10 rue A. Domon et Léonie Duquet, Paris, France

[4] Astrophysics Group, Cavendish Laboratory, University of Cambridge, J J Thomson Avenue, Cambridge CB3 0HE, U.K.

[5] Atacama Large Millimeter/submillimeter Array, ALMA Santiago Central Offices, Alonso de Cordova 3107, Vitacura, Casilla 763 0355, Santiago, Chile

[6] Australia Telescope National Facility, CSIRO, P.O. Box 76, Epping, NSW 1710, Australia

[7] CITA, University of Toronto, 60 St. George St., Toronto, ON M5S 3H8, Canada

[8] CNRS, IRAP, 9 Av. colonel Roche, BP 44346, F-31028 Toulouse cedex 4, France

[9] California Institute of Technology, Pasadena, California, U.S.A.

[10] Centre of Mathematics for Applications, University of Oslo, Blindern, Oslo, Norway

[11] DAMTP, University of Cambridge, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WA, U.K.

[12] DSM/Irfu/SPP, CEA-Saclay, F-91191 Gif-sur-Yvette Cedex, France

[13] DTU Space, National Space Institute, Juliane Mariesvej 30, Copenhagen, Denmark

[14] Departamento de Física, Universidad de Oviedo, Avda. Calvo Sotelo s/n, Oviedo, Spain

[15] Departamento de Matemáticas, Universidad de Oviedo, Avda. Calvo Sotelo s/n, Oviedo, Spain

[16] Department of Astronomy and Astrophysics, University of Toronto, 50 Saint George Street, Toronto, Ontario, Canada

[17] Department of Physics & Astronomy, The Open University, Milton Keynes, MK7 6AA, U.K.

[18] Department of Physics & Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver, British Columbia, Canada

[19] Department of Physics and Astronomy, University of Southern California, Los Angeles, California, U.S.A.

[20] Department of Physics, Gustaf Hällströmin katu 2a, University of Helsinki, Helsinki, Finland

[21] Department of Physics, Purdue University, 525 Northwestern Avenue, West Lafayette, Indiana, U.S.A.

[22] Department of Physics, University of California, Berkeley, California, U.S.A.

[23] Department of Physics, University of California, One Shields Avenue, Davis, California, U.S.A.

[24] Department of Physics, University of California, Santa Barbara, California, U.S.A.

[25] Department of Physics, University of Illinois at Urbana-Champaign, 1110 West Green Street, Urbana, Illinois, U.S.A.

[26] Dipartimento di Fisica G. Galilei, Università degli Studi di Padova, via Marzolo 8, 35131 Padova, Italy

[27] Dipartimento di Fisica, Università La Sapienza, P. le A. Moro 2, Roma, Italy

[28] Dipartimento di Fisica, Università degli Studi di Milano, Via Celoria, 16, Milano, Italy

[29] Dipartimento di Fisica, Università degli Studi di Trieste, via A. Valerio 2, Trieste, Italy

[30] Dipartimento di Fisica, Università di Ferrara, Via Saragat 1, 44122 Ferrara, Italy

[31] Dipartimento di Fisica, Università di Roma Tor Vergata, Via della Ricerca Scientifica, 1, Roma, Italy

[32] Discovery Center, Niels Bohr Institute, Blegdamsvej 17, Copenhagen, Denmark

[33] Dpto. Astrofísica, Universidad de La Laguna (ULL), E-38206 La Laguna, Tenerife, Spain

[34] European Southern Observatory, ESO Vitacura, Alonso de Cordova 3107, Vitacura, Casilla 19001, Santiago, Chile

[35] European Space Agency, ESAC, Planck Science Office, Camino bajo del Castillo, s/n, Urbanización Villafranca del

Castillo, Villanueva de la Cañada, Madrid, Spain

[36] European Space Agency, ESTEC, Keplerlaan 1, 2201 AZ Noordwijk, The Netherlands

[37] Haverford College Astronomy Department, 370 Lancaster Avenue, Haverford, Pennsylvania, U.S.A.

[38] Helsinki Institute of Physics, Gustaf Hällströmin katu 2, University of Helsinki, Helsinki, Finland

[39] IFSI/INAF, via del Fosso Cavaliere 100, 00133, Roma, Italy

[40] INAF - Osservatorio Astrofisico di Catania, Via S. Sofia 78, Catania, Italy

[41] INAF - Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, Padova, Italy

[42] INAF - Osservatorio Astronomico di Roma, via di Frascati 33, Monte Porzio Catone, Italy

[43] INAF - Osservatorio Astronomico di Trieste, Via G.B. Tiepolo 11, Trieste, Italy

[44] INAF/IASF Bologna, Via Gobetti 101, Bologna, Italy

[45] INAF/IASF Milano, Via E. Bassini 15, Milano, Italy

[46] INRIA, Laboratoire de Recherche en Informatique, Université Paris-Sud 11, Bâtiment 490, 91405 Orsay Cedex, France

[47] IPAG: Institut de Planétologie et d'Astrophysique de Grenoble, Université Joseph Fourier, Grenoble 1 / CNRS-INSU, UMR 5274, Grenoble, F-38041, France

[48] ISDC Data Centre for Astrophysics, University of Geneva, ch. d'Ecogia 16, Versoix, Switzerland

[49] Imperial College London, Astrophysics group, Blackett Laboratory, Prince Consort Road, London, SW7 2AZ, U.K.

[50] Infrared Processing and Analysis Center, California Institute of Technology, Pasadena, CA 91125, U.S.A.

[51] Institut d'Astrophysique Spatiale, CNRS (UMR8617) Université Paris-Sud 11, Bâtiment 121, Orsay, France

[52] Institut d'Astrophysique de Paris, CNRS UMR7095, Université Pierre & Marie Curie, 98 bis boulevard Arago, Paris, France

[53] Institute of Astronomy and Astrophysics, Academia Sinica, Taipei, Taiwan

[54] Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, U.K.

[55] Institute of Theoretical Astrophysics, University of Oslo, Blindern, Oslo, Norway

[56] Instituto de Astrofísica de Canarias, C/Vía Láctea s/n, La Laguna, Tenerife, Spain

[57] Instituto de Física de Cantabria (CSIC-Universidad de Cantabria), Avda. de los Castros s/n, Santander, Spain

[58] Istituto di Fisica del Plasma, CNR-ENEA-EURATOM Association, Via R. Cozzi 53, Milano, Italy

[59] Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, California, U.S.A.

[60] Jodrell Bank Centre for Astrophysics, Alan Turing Building, School of Physics and Astronomy, The University of Manchester, Oxford Road, Manchester, M13 9PL, U.K.

[61] Kavli Institute for Cosmology Cambridge, Madingley Road, Cambridge, CB3 0HA, U.K.

[62] LERMA, CNRS, Observatoire de Paris, 61 Avenue de l'Observatoire, Paris, France

[63] Laboratoire AIM, IRFU/Service d'Astrophysique - CEA/DSM - CNRS - Université Paris Diderot, Bât. 709, CEA-Saclay, F-91191 Gif-sur-Yvette Cedex, France

[64] Laboratoire Traitement et Communication de l'Information, CNRS (UMR 5141) and Télécom ParisTech, 46 rue Barrault F-75634 Paris Cedex 13, France

[65] Laboratoire de Physique Subatomique et de Cosmologie, CNRS, Université Joseph Fourier Grenoble I, 53 rue des

Martyrs, Grenoble, France

[66] Laboratoire de l'Accélérateur Linéaire, Université Paris-Sud 11, CNRS/IN2P3, Orsay, France

[67] Lawrence Berkeley National Laboratory, Berkeley, California, U.S.A.

[68] Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, 85741 Garching, Germany

[69] MilliLab, VTT Technical Research Centre of Finland, Tietotie 3, Espoo, Finland

[70] National University of Ireland, Department of Experimental Physics, Maynooth, Co. Kildare, Ireland

[71] Niels Bohr Institute, Blegdamsvej 17, Copenhagen, Denmark

[72] Observational Cosmology, Mail Stop 367-17, California Institute of Technology, Pasadena, CA, 91125, U.S.A.

[73] SISSA, Astrophysics Sector, via Bonomea 265, 34136, Trieste, Italy

[74] SUPA, Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, U.K.

[75] School of Physics and Astronomy, Cardiff University, Queens Buildings, The Parade, Cardiff, CF24 3AA, U.K.

[76] Space Sciences Laboratory, University of California, Berkeley, California, U.S.A.

[77] Spitzer Science Center, 1200 E. California Blvd., Pasadena, California, U.S.A.

[78] Stanford University, Dept of Physics, Varian Physics Bldg, 382 Via Pueblo Mall, Stanford, California, U.S.A.

[79] Université de Toulouse, UPS-OMP, IRAP, F-31028 Toulouse cedex 4, France

[80] University of Granada, Departamento de Física Teórica y del Cosmos, Facultad de Ciencias, Granada, Spain

[81] University of Miami, Knight Physics Building, 1320 Campo Sano Dr., Coral Gables, Florida, U.S.A.

[82] Warsaw University Observatory, Aleje Ujazdowskie 4, 00-478 Warszawa, Poland